


## E-Cadherin Antibody / CDH1 [clone CDH1/1525] (V3190)


Catalog No.	Formulation	Size
V3190-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3190-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3190SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	CDH1/1525
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P12830
<b>Localization</b>	Cytoplasmic, membranous
<b>Applications</b>	Immunofluorescence : 1-2ug/ml Western blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This E-Cadherin antibody is available for research use only.




Formalin-fixed, paraffin-embedded human colon carcinoma stained with E-Cadherin antibody (CDH1/1525).




Formalin-fixed, paraffin-embedded human colon carcinoma stained with E-Cadherin antibody (CDH1/1525).




Formalin-fixed, paraffin-embedded human prostate carcinoma stained with E-Cadherin antibody (CDH1/1525).




Formalin-fixed, paraffin-embedded human skin stained with E-Cadherin antibody (CDH1/1525).



Formalin-fixed, paraffin-embedded human placenta stained with E-Cadherin antibody (CDH1/1525).




Western blot analysis A) partial recombinant protein B) human stomach lysate using E-Cadherin antibody (CDH1/1525). Expected molecular weight: 135 kDa (precursor), 80-120 kDa (mature, depending on glycosylation level).




Immunofluorescent staining of human MCF7 cells with E-Cadherin antibody (clone CDH1/1525, green) and Reddot nuclear stain (red).

### Human Protein Microarray Specificity Validation




Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using E-Cadherin antibody (clone CDH1/1525). These results demonstrate the foremost specificity of the CDH1/1525 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free E-Cadherin antibody (clone CDH1/1525) as confirmation of integrity and purity.



Formalin-fixed, paraffin-embedded human breast adenocarcinoma stained with E-Cadherin antibody (CDH1/1525).

## Description

Recognizes a protein of 80-120kDa, identified as E-cadherin. Cadherins comprise a family of Ca<sup>2+</sup>-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH<sub>2</sub> terminal repeats. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as  $\beta$ -catenin, to regulate cadherin function. E-cadherin plays an important role in epithelial cell adhesion. A decreased expression of E-cadherin is associated with metastatic potential and poor prognosis in breast cancer, prostate and esophageal cancer. In combination with p120 Catenin, it is useful for the differentiation between ductal (E-cadherin +) and lobular (E-cadherin -) breast carcinomas. It may also help in diagnosis of mesothelioma.

## Application Notes

Optimal dilution of the E-Cadherin antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.

## Immunogen

A full length recombinant human protein was used as the immunogen for the E-Cadherin antibody.

## Storage

Store the E-Cadherin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).